




Contents lists available at ScienceDirect

JPRA Open

journal homepage: www.elsevier.com/locate/jpra

Review Article

The evolution of perforator flaps and the future of microsurgery

Joon Pio Hong^{a,*}, Jin Geun Kwon^a, Hyunsuk Peter Suh^a,
Changsik John Pak^a, Hyung Bae Kim^a, Hyun Ho Han^a,
Hyungjoo Noh^b, Jae Young Hur^c, Erin Brown^d

^a Department of Plastic Surgery, Asan Medical Center, University of Ulsan College of Medicine, Seoul, Korea^b Department of Plastic and Reconstructive Surgery, Yas Clinic Khalifa City, Abu Dhabi, United Arab Emirates^c Plastic Surgery, Young Young Plastic Surgery Clinic, Seoul, Korea^d Department of Plastic and Reconstructive Surgery, University of British Columbia, Vancouver, Canada

ARTICLE INFO

Article history:

Received 10 September 2025

Accepted 11 September 2025

Available online xxx

Keywords:

Microsurgery

Supermicrosurgery

Perforator flap

Digital microscope

Robotic microsurgery platform

ABSTRACT

Perforator flaps have transformed reconstructive microsurgery by enabling tissue-specific reconstruction while preserving muscle, fascia, and nerves, minimizing donor site morbidity. Modern techniques—including superthin, ultrathin, and pure skin flaps—enhance flap precision, safety, and versatility. Imaging tools such as CT angiography and high-resolution ultrasound allow accurate mapping of perforator anatomy, improving flap design and outcomes. Challenges like short pedicle length and flap thickness are addressed through perforator-to-perforator supermicrosurgery, enabling anastomosis of submillimeter vessels with minimal disruption. Advances in high-magnification microscopes, ultrafine microsutures, robotic platforms, and digital exoscopes further expand surgical capabilities, improve ergonomics, and shorten the learning curve. Looking ahead, artificial intelligence and augmented reality promise to automate microsurgical tasks, enhance visualization, and optimize functional and aesthetic results. Collectively, these innovations are pushing reconstructive microsurgery toward the

* Corresponding author.

E-mail address: joonphong@amc.seoul.kr (J.P. Hong).

<https://doi.org/10.1016/j.jptra.2025.09.016>

2352-5878/© 2025 The Author(s). Published by Elsevier Ltd on behalf of British Association of Plastic, Reconstructive and Aesthetic Surgeons. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Please cite this article as: J.P. Hong, J.G. Kwon, H.P. Suh et al., The evolution of perforator flaps and the future of microsurgery, JPRA Open, <https://doi.org/10.1016/j.jptra.2025.09.016>

"reconstructive elevator" ideal, achieving safer, more efficient, and highly customized outcomes for patients.

© 2025 The Author(s). Published by Elsevier Ltd on behalf of British Association of Plastic, Reconstructive and Aesthetic Surgeons. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Introduction of perforator flaps

In the era of the reconstructive elevator concept, achieving both functional and aesthetic surgical outcomes with maximal efficiency has become the standard (Figure 1).¹ However, the development of this approach was gradual. In the early days of reconstructive surgery, the primary focus was on wound closure, often without detailed understanding of flap vascularity. The importance of vascular knowledge became evident, as random-pattern flaps with unpredictable blood supply were limited in their ability to cover larger defects. The introduction of axial-pattern flaps, with defined blood supply to muscle and musculocutaneous tissue, provided a framework for flap design. Nevertheless, these flaps often required sacrificing muscle, resulting in functional donor site morbidity and undesirable bulk. In 1989, Koshima and Soeda first articulated the concept of perforator flaps, which have since gained popularity and are now considered a cornerstone of modern reconstructive microsurgery.^{2,3} The concept of perforator flaps—utilizing autologous tissue while preserving fascia, muscle, and nerves to minimize donor site morbidity—was revolutionary, offering an early glimpse into the future of flap surgery.^{3,4} The most frequently used perforator flaps nowadays are the deep inferior epigastric perforator (DIEP) flap, superior and inferior gluteal (SGAP/IGAP) flap mainly for breast reconstruction, anterolateral thigh (ALT) flap and the thoracodorsal artery perforator (TDAP) flap mainly for the ex-




Figure 1. The reconstructive elevator concept. Unlike the ladder's sequential steps, the elevator allows surgeons to skip rungs and directly access more advanced or appropriate solutions, acknowledging that technological advances have made complex options more viable.

tremities and the head and neck region.^{5–10} These perforator flaps have allowed to reconstruct the defect with the most similar tissue possible truly with minimal donor site morbidity.

Additional advantages of perforator flaps

In addition to the features mentioned above such as providing like-with-like reconstruction for resurfacing, minimizing donor site morbidity by preserving muscle function, additional advantages such as being able to provide sensation, provide long pedicles in selected flaps, ability to design versatile flaps such as chimeric style flaps, and to be able to customize according to the defect are well documented.^{4,11–17} The perforator flap principle can also be used in a local flap reflecting the same versatility as being used as a free flap.^{18–25}

A study by Hong et al. shows that even if the perforator flaps are not innervated, 72 % protective sensation, 21 % fine tactile, 49 % temperature, and 64 % pain peripheral sensation were present at 1 year after the reconstruction and continued to improve at 34 months.¹² The sensory recovery occurs from periphery to the central over time. Understanding the fate of sensory recovery for non-innervated perforator flaps will allow surgeons to further customize the reconstruction based on the defect and maximize efficiency, moving a step closer toward the ideal elevator approach.

A chimeric flap is constructed from multiple, distinct tissue components, such as skin, muscle, or bone, that all have independent vascular supplies but share a single, common parent blood vessel.^{2,23,24} This surgical technique utilizing the vascular anatomy of perforator flaps allows surgeons to reconstruct complex defects with a single vascular pedicle, combining different tissue types to match the specific requirements of the defect site again allowing to move closer to the elevator approach.²⁶

The perforator flaps provide many advantages that can be considered when selecting the flap. An algorithmic thinking allows to select the right perforator flap for different defects.^{25,27} Perforator flap can be chosen based on the requirements such as pedicle length, thickness of flap, need for chimeric flap, position of the patient on the surgical table and other factors. Understanding the characteristics of the selected perforator flap allows to better reconstruct the defect.

Overcoming the limitations of perforator flaps

As quoted by Dr Fu-Chan Wei, the ALT flap has been acknowledged as an ideal flap for reconstruction due to its versatility, reliable vascular anatomy, long vascular pedicle, and minimal donor site morbidity.⁹ It offers supple, durable coverage that can be tailored in size and shape for various defects across the body, from the head to the foot. Its reliable blood supply facilitates microvascular




Figure 2. Various planes for elevating the perforator flap.

anastomosis, while the option to include the lateral femoral cutaneous nerve allows for sensate flaps, enhancing functional outcome.^{28,29} The ability to harvest it as a two-team approach also reduces operating time, further supporting its role as a workhorse flap. In most of the perforator flap, these advantages apply to some extent as well. However, with increasing knowledge of perforator flaps and advancement of techniques, the standard for the ideal flap is steadily rising. Some may say that the linear scar on the thigh or the skin graft of the donor site after the harvest may not be so aesthetically favorable. Also in relatively obese patients, the flap may be too thick to accommodate the recipient site often requiring a second stage debulking of the flap. Thus the authors searched for an inherently thinner flap such as superficial circumflex iliac artery perforator (SCIP) flap originally introduced by Dr Koshima and medial sural artery perforator (MSAP) flap introduced by Dr Cavadas to resurface shallow defects.^{30–40} However, these flaps also have their limitations such as having short pedicle or limited skin dimension. Thus, various methods for thinning were introduced to avoid inaccurate reconstructions such as skin grafts on the flap margin and to over-sized skin flap to accommodate the access thickness of the flap. Although intraoperative debulking procedures are widely practiced, there is always the question of how much thinning is possible without injuring the flap vascularity.^{41–44} Identifying a safe plane of elevation while preserving the vascular anatomy and flow may be the ideal approach to obtain thin flaps. Widely utilized planes of elevation are subfascial (harvesting the flap just below the deep fascia; including the deep fascia within the flap) and suprafascial (elevation is above the deep fascia) frequently still being far from obtaining a thin flap especially in obese patients. In order to safely elevate a thin flap, the authors introduced identifying and elevating the flap on the superficial fascia plane between the deep fat and the superficial fat and named it as “superthin flap.”^{8,45–54} New planes have subsequently been introduced such as the “ultrathin” which is elevating within the superficial fat and “pure skin” which is elevating with minimal fat attached to the flap obtaining even thinner flaps (Figure 2).^{54–62} Thus the issue with bulky or thick flaps has now become less of an concern and these elevation techniques on various planes contributed a step closer to the elevator approach.⁵⁴ Furthermore, understanding the perforator and its pathway allows not only to take the flap as a pure skin flap but also allows to take a single component of the flap such as pure fat flap.⁶³

A key factor in elevating thin flaps is understanding the vascular anatomy, particularly the course of the perforators. Knowing this pathway makes flap elevation both safer and more efficient. While




Figure 3. The ultrahigh high frequency ultrasound showing the point where the perforator penetrates the deep fascia (shown in blue in the color Doppler mode). The superficial fascia as well as the deep fascia is clearly shown.

the source vessel anatomy in perforator flaps is usually well defined, the intramuscular or subcutaneous course of the perforators can often be complex and difficult to trace. For example, in ALT flaps, microsurgeons generally know the location of the main perforator, which lies at the mid-point between the anterior superior iliac spine and the medial femoral head. However, the course of the perforator through muscle or subcutaneous fat is difficult to determine without proper imaging. With adequate radiologic evaluation, surgeons can map the perforator pathway to improve flap design and better understand its intramuscular course, which makes dissection easier. CT angiography is often used to provide this information.^{64–68} As the elevation technique advanced to involve multiple layers, understanding the perforator pathway within the subcutaneous fat became essential for preserving skin vascularity. With ultrasound (Aplio i800, Canon medical systems, Tokyo, Japan), surgeons can now visualize not only the real-time course of these vessels but also their physiological characteristics, including velocity, flow volume, tissue perfusion and more (Figure 3).^{20,52,69–72} The authors believe that ultrasound helps surgeons work from different planes, design chimera flaps, and understand flap blood flow, which together improve the accuracy and precision of flap design.

Opening the era of perforator flaps mixed with the concept of freestyle, the choices for perforator flaps increased immensely. Basically, any perforator can be used as a pedicle for a skin flap.^{15,19,20} The challenge, however, was the pedicle length. A typical example can be the SCIP or a posterior interosseous artery perforator (PIAP) flap. Although seen as an inherently a thin flap, often the pedicle can be quite short without modification.^{14,40,73} To overcome the challenges of short pedicle length, the authors introduced and advocated the perforator-to-perforator approach using the supermicrosurgery concept.^{74–76} This approach allow to preserve main larger arteries without any disruption, avoids donor site morbidity by taking only a short segment of the perforator avoiding extensive dissection, increases donor site options, may provide physiologic inflow and shortens operation time opening to door to minimal invasive reconstruction.^{74,76–78}

Various new innovations continue in the field of perforator flaps to not only overcome challenges intrinsic from using perforator flaps but also to increase the precision, accuracy and ultimately allowing a step further to a true elevator approach. One might question whether these innovations are merely visions of the future, distant from current surgical practice.^{79,80} Yet, surgical progress has always relied on pioneers who introduce and validate new concepts, while others adopt them more




Figure 4. From the top right, that has an arrow with 0.03 mm (30 microns), it is 12–0 s, followed by 12–0, 11–0, and 10–0 underneath each needle. The needle that has 0.12 mm (120 μ m) is a 9–0 microsuture. To show the relative size, a cardiovascular needle is shown on the lower part of the photo above the ruler. Image courtesy of Crownjun.

gradually due to variations in resources, training, or institutional readiness.^{81–83} It remains essential, however, for followers to remain engaged with innovation, as timely adoption allows patients to benefit from safer, more efficient, and more effective treatments.

The future of microsurgery

The introduction and widespread adoption of perforator flaps created the demand for higher magnification, ultrafine microsurgical instruments, finer microsutures for small-caliber vessels, and advanced imaging modalities. Technology adapted from space telescopes has been integrated into surgical microscopes, allowing magnification of up to 77 times (MM51, Mitaka, Tokyo, Japan). Furthermore, microsutures with needles as small as 0.8 mm in length and 30 μm in diameter (12–0, Crown-jun, Tokyo, Japan) have enabled anastomosis of vessels as narrow as 0.4 mm in diameter (Figure 4). Collectively, these advances have not only made perforator-to-perforator supermicrosurgery a reality but have also established supermicrosurgery as a distinct class of reconstructive surgery.⁸⁴ Many surgeons question how steep the learning curve is before one can reliably handle vessels of such small dimensions. The introduction of robotic microsurgical platforms (Symani, MMI, Jacksonville, Florida, USA) has addressed this challenge by providing surgeons—even those with modest microsurgical experience—with enhanced precision and dexterity, tremor elimination, improved ergonomics, and reduced fatigue (Figure 5).^{85–87} Artificial intelligence is expected to play a key role in the future automation of microsurgery. In this context, digital microscopes and exoscopes are being introduced (ScopeEye, MedithinQ, Seoul, Korea) with robust capabilities for data collection (Figure 6). These systems provide surgeons with three-dimensional digital visualization of the operative field, with the potential to incorporate augmented reality overlays and enable real-time communication with peers during surgery. Such innovations aim to enhance precision, safety, and visualization for complex tasks, while also improving ergonomics for the surgeon.



Figure 5. The robotic microsurgery platform Symani. Note the microneedle holder and forceps attached to the robotic wrist in which the master (surgeon) controls from the console. Image courtesy of Medical Microinstruments, Inc.



Figure 6. A surgeon wearing a digital visor (Scope Eye) is shown. The visor imports the image from the versatile microscope which has a small foot print. The assistant is also seen wearing the visor together performing a lymphovenous anastomosis for a 0.4 mm caliber. The same image is also transported to the large flat screen which can be seen in 3 dimension using a 3D glass.

Conclusion

Perforator flaps have revolutionized reconstructive microsurgery by combining versatility with minimal donor site morbidity. Advances in imaging, flap elevation techniques, and supermicrosurgery have enhanced precision, safety, and aesthetic outcomes. Robotic platforms, digital exoscopes, and emerging AI technologies promise to further improve efficiency, visualization, and surgical ergonomics. Together, these innovations are driving the field toward the “reconstructive elevator” ideal—maximizing functional restoration, aesthetics, and patient-centered results.

Declaration of competing interest

The authors have no commercial associations or financial disclosures to declare.

References

- Gottlieb LJ, Krieger LM. From the reconstructive ladder to the reconstructive elevator. *Plast Reconstr Surg.* 1994;93:1503–1504.
- Koshima I, Soeda S. Inferior epigastric artery skin flaps without rectus abdominis muscle. *Br J Plast Surg.* 1989;42:645–648.
- Hallock GG. Evaluation of fasciocutaneous perforators using color duplex imaging. *Plast Reconstr Surg.* 1994;94:644–651.
- Wei FC, Celik N. Perforator flap entity. *Clin Plast Surg.* 2003;30:325–329.
- Lee SY, Seok MC, Park BY. Surgical considerations of one-stage reconstruction of large extremity defects using a thin deep inferior epigastric perforator flap. *Arch Plast Surg.* 2023;50:586–592.
- Kim JS, Lee HH, Koh SH, et al. Hand reconstruction using anterolateral thigh free flap by terminal perforator-to-digital artery anastomosis: retrospective analysis. *Arch Plast Surg.* 2024;51:87–93.
- Speck NE, Grufman V, Farhadi J. Trends and innovations in autologous breast reconstruction. *Arch Plast Surg.* 2023;50:240–247.

8. Hong JP, Yim JH, Malzone G, et al. The thin gluteal artery perforator free flap to resurface the posterior aspect of the leg and foot. *Plast Reconstr Surg.* 2014;133:1184–1191.
9. Wei FC, Jain V, Celik N, et al. Have we found an ideal soft-tissue flap? An experience with 672 anterolateral thigh flaps. *Plast Reconstr Surg.* 2002;109:2219–2226 discussion 27–30.
10. Koshima I, Narushima M, Mihara M, et al. New thoracodorsal artery perforator (TAPcp) flap with capillary perforators for reconstruction of upper limb. *J Plast Reconstr Aesthet Surg.* 2010;63:140–145.
11. Jaiswal D, Saxena B, Mathews S, et al. TFL perforator flap complementing and completing the ALT-AMT flap axis. *Arch Plast Surg.* 2024;51:408–416.
12. Kim HB, Cho JM, Abdelfattah U, et al. The fate of sensation in noninnervated perforator free flaps in the lower extremity. *Plast Reconstr Surg.* 2023;152:440–449.
13. Kwon JG, Cho MJ, Pak CJ, Suh HP, Hong JP. A retrospective case series on free flap reconstruction for ischemic diabetic foot: the nutrient flap further explained. *Plast Reconstr Surg.* 2022;149:1452–1461.
14. Kwon JG, Pereira N, Tonaree W, et al. Long pedicled superficial circumflex iliac artery flap based on a medial superficial branch. *Plast Reconstr Surg.* 2021;148:615e–619e.
15. Wei FC, Mardini S. Free-style free flaps. *Plast Reconstr Surg.* 2004;114:910–916.
16. Wallace CG, Kao HK, Jeng SF, Wei FC. Free-style flaps: a further step forward for perforator flap surgery. *Plast Reconstr Surg.* 2009;124:e419–e426.
17. Song JW, Ben-Nakhi M, Hong JP. Reconstruction of lower extremity with perforator free flaps by free style approach in pediatric patients. *J Reconstr Microsurg.* 2012;28:589–594.
18. Hong JP, Choi D, Lee Y, et al. Using the gluteal artery perforator flap to reconstruct sacral sore. *Plast Reconstr Surg Glob Open.* 2017;5:e1368.
19. Park SW, Oh TS, Eom JS, et al. Freestyle multiple propeller flap reconstruction (jigsaw puzzle approach) for complicated back defects. *J Reconstr Microsurg.* 2015;31:261–267.
20. Kim HB, Han SJ, Hong JP, Suh HP. The preoperative planning, design, and execution of the freestyle propeller flap: a detailed description and the case series. *Arch Plast Surg.* 2025;52:239–246.
21. Datli A, Suh H, Kim YC, Choi DH, Hong JP. Free-style deepithelialized propeller flaps: an ideal local flap to obliterate wounds with dead space. *Plast Reconstr Surg Glob Open.* 2017;5:e1249.
22. Kedar DJ, Pak CJ, Suh HP, Hong JP. Propeller flaps in the posterior trunk. *Semin Plast Surg.* 2020;34:176–183.
23. Hallock GG. The combined parascapular fasciocutaneous and latissimus dorsi muscle conjoined free flap. *Plast Reconstr Surg.* 2008;121:101–107.
24. Koshima I, Yamamoto H, Hosoda M, et al. Free combined composite flaps using the lateral circumflex femoral system for repair of massive defects of the head and neck regions: an introduction to the chimeric flap principle. *Plast Reconstr Surg.* 1993;92:411–420.
25. Abdelfattah U, Power HA, Song S, et al. Algorithm for free perforator flap selection in lower extremity reconstruction based on 563 cases. *Plast Reconstr Surg.* 2019;144:1202–1213.
26. Hallock GG. The chimeric propeller flap. *Semin Plast Surg.* 2020;34:207–209.
27. Hallock GG. An overview algorithm for perforator free flap coverage of all zones of the mangled upper extremity. *J Hand Surg Glob Online.* 2025;7:277–283.
28. Hong JP. Reconstruction of the diabetic foot using the anterolateral thigh perforator flap. *Plast Reconstr Surg.* 2006;117:1599–1608.
29. Hong JP, Kim EK. Sole reconstruction using anterolateral thigh perforator free flaps. *Plast Reconstr Surg.* 2007;119:186–193.
30. Choi JW, Nam SY, Choi SH, et al. Applications of medial sural perforator free flap for head and neck reconstructions. *J Reconstr Microsurg.* 2013;29:437–442.
31. Cavadas PC, Sanz-Gimenez-Rico JR, Gutierrez-de la Camara A, et al. The medial sural artery perforator free flap. *Plast Reconstr Surg.* 2001;108:1609–1615 discussion 16–7.
32. Koshima I, Nanba Y, Tsutsui T, et al. Superficial circumflex iliac artery perforator flap for reconstruction of limb defects. *Plast Reconstr Surg.* 2004;113:233–240.
33. Sapino G, Osinga R, Maruccia M, et al. Role of the gastrocnemius musculocutaneous with a propeller style skin flap in knee region reconstruction: indications and pitfalls. *Arch Plast Surg.* 2023;50:593–600.
34. Suh HS, Jeong HH, Choi DH, Hong JP. Study of the medial superficial perforator of the superficial circumflex iliac artery perforator flap using computed tomographic angiography and surgical anatomy in 142 patients. *Plast Reconstr Surg.* 2017;139:738–748.
35. Choi DH, Goh T, Cho JY, Hong JP. Thin superficial circumflex iliac artery perforator flap and supermicrosurgery technique for face reconstruction. *J Craniofac Surg.* 2014;25:2130–2133.
36. Goh TL, Park SW, Cho JY, Choi JW, Hong JP. The search for the ideal thin skin flap: superficial circumflex iliac artery perforator flap—a review of 210 cases. *Plast Reconstr Surg.* 2015;135:592–601.
37. Hong JP. The superficial circumflex iliac artery perforator flap in lower extremity reconstruction. *Clin Plast Surg.* 2021;48:225–233.
38. Hong JP, Sun SH, Ben-Nakhi M. Modified superficial circumflex iliac artery perforator flap and supermicrosurgery technique for lower extremity reconstruction: a new approach for moderate-sized defects. *Ann Plast Surg.* 2013;71:380–383.
39. Suh YC, Hong JP, Suh HP. Elevation technique for medial branch based superficial circumflex iliac artery perforator flap. *Handchir Mikrochir Plast Chir.* 2018;50:256–258.
40. Sisternas Hernandez L, Lopez Fernandez S, Zamora Alarcon PD, et al. Posttraumatic scrotal reconstruction with a pedicled “extended” superficial circumflex iliac artery perforator flap: a case report. *Arch Plast Surg.* 2024;51:327–331.
41. Ross GL, Dunn R, Kirkpatrick J, et al. To thin or not to thin: the use of the anterolateral thigh flap in the reconstruction of intraoral defects. *Br J Plast Surg.* 2003;56:409–413.
42. Sharabi SE, Hafez DA, Koshy JC, et al. Is primary thinning of the anterolateral thigh flap recommended? *Ann Plast Surg.* 2010;65:555–559.
43. Kimura N. A microdissected thin tensor fasciae latae perforator flap. *Plast Reconstr Surg.* 2002;109:69–77 discussion 78–80.
44. Park BY. Flap thinning: defatting after conventional elevation. *Arch Plast Surg.* 2018;45:314–318.

45. Hong JP, Choi DH, Suh H, et al. A new plane of elevation: the superficial fascial plane for perforator flap elevation. *J Reconstr Microsurg.* 2014;30:491–496.
46. Hong JP, Chung IW. The superficial fascia as a new plane of elevation for anterolateral thigh flaps. *Ann Plast Surg.* 2013;70:192–195.
47. Jeong HH, Hong JP, Suh HS. Thin elevation: a technique for achieving thin perforator flaps. *Arch Plast Surg.* 2018;45:304–313.
48. Sakarya AH, Do N. Nomenclature of thin and super-thin flaps-comment on: outcomes of subfascial, suprafascial, and super-thin anterolateral thigh flaps: tailoring thickness without added morbidity. *J Reconstr Microsurg.* 2018;34:e3–e4.
49. Park SO, Chang H, Imanishi N. Anatomic basis for flap thinning. *Arch Plast Surg.* 2018;45:298–303.
50. Kim KN, Hong JP, Park CR, Yoon CS. Modification of the elevation plane and defatting technique to create a thin thoracodorsal artery perforator flap. *J Reconstr Microsurg.* 2016;32:142–146.
51. Suh YC, Kim NR, Jun DW, Lee JH, Kim YJ. The perforator-centralizing technique for super-thin anterolateral thigh perforator flaps: minimizing the partial necrosis rate. *Arch Plast Surg.* 2021;48:121–126.
52. Visconti G, Bianchi A, Hayashi A, et al. Thin and superthin perforator flap elevation based on preoperative planning with ultrahigh-frequency ultrasound. *Arch Plast Surg.* 2020;47:365–370.
53. Cha HG, Hur J, Ahn C, Hong JP, Suh HP. Ultrathin anterolateral thigh free flap: an adipocutaneous flap with the most superficial elevation plane. *Plast Reconstr Surg.* 2023;152:718e–723e.
54. Kwon JG, Brown E, Suh HP, Pak CJ, Hong JP. Planes for perforator/skin flap elevation-definition, classification, and techniques. *J Reconstr Microsurg.* 2023;39:179–186.
55. Narushima M, Yamasoba T, Iida T, et al. Pure skin perforator flaps: the anatomical vascularity of the superthin flap. *Plast Reconstr Surg.* 2018;142:351e–360e.
56. Kimura N, Satoh K. Consideration of a thin flap as an entity and clinical applications of the thin anterolateral thigh flap. *Plast Reconstr Surg.* 1996;97:985–992.
57. Visconti G, Bianchi A, Hayashi A, Salgarello M. Pure skin perforator flap direct elevation above the subdermal plane using preoperative ultra-high frequency ultrasound planning: a proof of concept. *J Plast Reconstr Aesthet Surg.* 2019;72:1700–1738.
58. Bhadkamkar MA, Wolfswinkel EM, Hatf DA, et al. The ultra-thin, fascia-only anterolateral thigh flap. *J Reconstr Microsurg.* 2014;30:599–606.
59. Patel KM, Shauly O, Gould DJ. Introducing the subdermal free flap: preserving the ultrathin-free skin flap option in morbidly obese patients. *J Surg Oncol.* 2018;118:403–406.
60. Yamamoto T, Yoshimatsu H, Kikuchi K. Free prepuce perforator flap: ultra-thin superficial penile artery perforator flap for a dorsal finger defect. *Microsurgery.* 2017;37:252–255.
61. Narushima M, Yamasoba T, Iida T, et al. Pure skin perforator flap for microtia and congenital aural atresia using supermicrosurgical techniques. *J Plast Reconstr Aesthet Surg.* 2011;64:1580–1584.
62. Yamamoto T, Yamamoto N, Fuse Y, et al. Subdermal dissection for elevation of pure skin perforator flaps and superthin flaps: the dermis as a landmark for the most superficial dissection plane. *Plast Reconstr Surg.* 2021;147:470–478.
63. Lee SJ, Cho J, Pak C, Suh H, Hong JP. “Pure fat flap”—perforator-based adiposal layer only flap for lateral ankle reconstruction. *Arch Plast Surg.* 2025;52:41–45.
64. Kim EK, Kang BS, Hong JP. The distribution of the perforators in the anterolateral thigh and the utility of multidetector row computed tomography angiography in preoperative planning. *Ann Plast Surg.* 2010;65:155–160.
65. Masia J, Clavero JA, Larranaga JR, et al. Multidetector-row computed tomography in the planning of abdominal perforator flaps. *J Plast Reconstr Aesthet Surg.* 2006;59:594–599.
66. Masia J, Clavero JA, Larranaga J, Vives L, Pons G. Preoperative planning of the abdominal perforator flap with multidetector row computed tomography: 3 years of experience. *Plast Reconstr Surg.* 2008;122:80e–81e.
67. Rozen WM, Ashton MW, Stella DL, et al. Developments in perforator imaging for the anterolateral thigh flap: CT angiography and CT-guided stereotaxy. *Microsurgery.* 2008;28:227–232.
68. Rozen WM, Phillips TJ, Ashton MW, et al. Preoperative imaging for DIEA perforator flaps: a comparative study of computed tomographic angiography and Doppler ultrasound. *Plast Reconstr Surg.* 2008;121:9–16.
69. Cho MJ, Kwon JG, Pak CJ, Suh HP, Hong JP. The role of duplex ultrasound in microsurgical reconstruction: review and technical considerations. *J Reconstr Microsurg.* 2020;36:514–521.
70. Kehrer A, Sachanadani NS, da Silva NPB, et al. Step-by-step guide to ultrasound-based design of alt flaps by the microsurgeon—basic and advanced applications and device settings. *J Plast Reconstr Aesthet Surg.* 2020;73:1081–1090.
71. Hong JP, Hur J, Kim HB, Park CJ, Suh HP. The use of color duplex ultrasound for local perforator flaps in the extremity. *J Reconstr Microsurg.* 2022;38:233–237.
72. Hong JP. The Color Duplex ultrasound: the reconstructive surgeon's stethoscope. *J Reconstr Microsurg.* 2022;38:169.
73. Yoon CS, Noh HJ, Malzone G, et al. Posterior interosseous artery perforator-free flap: treating intermediate-size hand and foot defects. *J Plast Reconstr Aesthet Surg.* 2014;67:808–814.
74. Hong JP, Koshima I. Using perforators as recipient vessels (supermicrosurgery) for free flap reconstruction of the knee region. *Ann Plast Surg.* 2010;64:291–293.
75. Kim KN, Hong JP, Park SW, Kim SW, Yoon CS. Overcoming the obstacles of the ilizarov device in extremity reconstruction: usefulness of the perforator as the recipient vessel. *J Reconstr Microsurg.* 2015;31:420–425.
76. Power HA, Cho J, Kwon JG, et al. Are perforators reliable as recipient arteries in lower extremity reconstruction? Analysis of 423 free perforator flaps. *Plast Reconstr Surg.* 2022;149:750–760.
77. Basaran K, Datli A, Sagir M, Sar M, Bilgi E. Blunt trauma induced closed femoral bone fracture in a rat model: are vessels safe to use for microsurgery? Further insight into the zone of injury concept. *J Reconstr Microsurg.* 2025;41:149–155.
78. Hong JP, Kim HB, Park CJ, Suh HP. Using Duplex ultrasound for recipient vessel selection. *J Reconstr Microsurg.* 2022;38:200–205.
79. Datli A, Karasoy I, Genc Y, Pilanci O. Challenges of setting up a lower extremity reconstruction practice in a constrained environment. *J Reconstr Microsurg.* 2021;37:67–74.
80. Ferreira N, Sabharwal S, Hosny GA, et al. Limb reconstruction in a resource-limited environment. *SICOT J.* 2021;7:66.
81. Hong JP. Innovation in plastic surgery-why and how? *Arch Plast Surg.* 2021;48:471–472.
82. Hallock GG. Innovations: a dawning of a new age. *Arch Plast Surg.* 2021;48:147–148.

83. Neligan PC. Innovation and collaboration in plastic surgery. *Arch Plast Surg.* 2024;51:147–149.
84. Hong JPJ, Song S, Suh HSP. Supermicrosurgery: principles and applications. *J Surg Oncol.* 2018;118:832–839.
85. Cho J, Kim D, Kim T, et al. Further validating the robotic microsurgery platform through preclinical studies on rat femoral artery and vein. *J Reconstr Microsurg.* 2025;41:595–605.
86. Innocenti M. Back to the future: robotic microsurgery. *Arch Plast Surg.* 2022;49:287–288.
87. Barbon C, Grunherz L, Uyulmaz S, Giovanoli P, Lindenblatt N. Exploring the learning curve of a new robotic microsurgical system for microsurgery. *JPRAS Open.* 2022;34:126–133.